

EMERGING INFECTIOUS DISEASES IN WILDLIFE: ZONOTIC RISKS, ECOLOGY AND CONTROL STRATEGIES

DILEEPU KUMAR BANISETTI

Assistant Professor, Department of Botany, Maharajah's College (Autonomous),
Affiliated to Andhra University, Vizayanagaram.

Article History: Received: 11 Oct 2025, Revised: 22 Nov 2025, Accepted: 04 Dec 2025

*Corresponding Author

Dr. Dileepu Kumar Baniseti


Abstract

Emerging infectious diseases (EIDs) originating from wildlife represent a growing global threat to public health, biodiversity, and ecosystem stability. The majority of recently emerging human pathogens are zoonotic, with wildlife acting as natural reservoirs. Anthropogenic pressures such as habitat destruction, climate change, wildlife trade, and agricultural expansion have intensified human-wildlife interactions, facilitating pathogen spillover. The ecology of wildlife diseases is shaped by host diversity, pathogen evolution, environmental conditions, and socio-economic factors. This review examines the role of wildlife in zoonotic disease emergence, key ecological drivers, mechanisms of spillover, and current control and mitigation strategies. Emphasis is placed on surveillance, One Health approaches, habitat management, and policy interventions to reduce future zoonotic risks while conserving wildlife populations.

Keywords: *Emerging infectious diseases; Wildlife; Zoonoses; Disease ecology; Spillover; One Health; Control strategies.*

This article is licensed under a Creative Commons Attribution-Non-commercial 4.0 International License.

Copyright © 2025 Author(s) retains the copyright of this article.



INTRODUCTION

Emerging infectious diseases are defined as infections that have newly appeared in a population or are rapidly increasing in incidence or geographic distribution. Over the past few decades, EIDs have increased significantly, with more than 60% of human infectious diseases being zoonotic and approximately 70% of these originating from wildlife reservoirs [1]. Wildlife species harbor a vast diversity of pathogens, many of which have the potential to infect humans under favorable ecological and socio-environmental conditions [2]. Understanding the ecological basis of disease emergence in wildlife is therefore critical for predicting and preventing future zoonotic outbreaks.

WILDLIFE AS RESERVOIRS OF ZONOTIC PATHOGENS

Wildlife species serve as natural reservoirs for a wide range of pathogens, including viruses, bacteria, protozoa, and helminths. Bats and rodents are particularly important reservoirs due to their species richness, wide geographic distribution, and ability to coexist with humans [3]. Examples include coronaviruses and henipaviruses in bats, hantaviruses in rodents, and avian influenza viruses in wild birds [4]. These hosts often exhibit limited clinical symptoms, allowing long-term pathogen persistence and transmission within populations [5].

ECOLOGICAL DRIVERS OF DISEASE EMERGENCE

1. Land Use Change and Habitat Fragmentation

Deforestation, agricultural expansion, mining, and urbanization disrupt natural ecosystems and increase contact between wildlife, livestock, and humans. Such changes can alter host density, vector abundance, and pathogen transmission dynamics, increasing spillover risk [6].

2. Climate Change

Climate change influences the distribution and seasonal activity of hosts, vectors, and pathogens. Rising temperatures and altered precipitation patterns have expanded the geographic range of vectors such as mosquitoes and ticks, facilitating the spread of vector-borne zoonoses [7].

3. Wildlife Trade and Consumption

Legal and illegal wildlife trade, including live animal markets and bushmeat consumption, creates high-risk interfaces for cross-species pathogen transmission. These practices have been implicated in the emergence of several zoonotic diseases, including SARS and Ebola virus disease [8].

4. Biodiversity Loss

Biodiversity loss can increase disease risk by favoring competent reservoir hosts, a phenomenon known as the "dilution effect." Reduced species diversity may

enhance pathogen transmission by increasing encounters between hosts and vectors [9].

MECHANISMS OF ZONOTIC SPILLOVER

Spillover occurs when a pathogen maintained in wildlife successfully infects humans. This process depends on pathogen prevalence in reservoir hosts, shedding intensity, environmental survival, and frequency of human exposure [10]. Human behaviors such as hunting, farming, and encroachment into wildlife habitats play a crucial role in facilitating spillover events.

SURVEILLANCE AND DISEASE ECOLOGY RESEARCH

Wildlife disease surveillance is essential for early detection of emerging pathogens. Integrated surveillance systems combining field sampling, molecular diagnostics, ecological modeling, and community-based reporting have proven effective in identifying high-risk regions and hosts [11]. Advances in genomic sequencing have further improved understanding of pathogen evolution and transmission pathways.

CONTROL AND MITIGATION STRATEGIES

1. One Health Approach

The One Health framework recognizes the interconnectedness of human, animal, and environmental health. Coordinated efforts across disciplines improve surveillance, risk assessment, and outbreak response for zoonotic diseases [12].

2. Habitat Conservation and Management

Protecting intact ecosystems and minimizing habitat fragmentation reduce forced interactions between wildlife and humans, thereby lowering spillover risk [6].

3. Wildlife Vaccination and Population Management

Targeted vaccination programs, such as oral rabies vaccination in wild carnivores, have successfully reduced disease transmission. However, ecological impacts and feasibility must be carefully evaluated [13].

4. Policy and Regulation

Strengthening regulations on wildlife trade, enforcing biosecurity measures, and improving public awareness are essential components of zoonotic disease prevention [14].

CASE STUDIES

1. Ebola Virus Disease

Ebola virus spillover has been linked to contact with infected wildlife, particularly bats and non-human primates. Subsequent human-to-human transmission has caused severe outbreaks in Africa, highlighting the importance of wildlife surveillance and community engagement [15].

2. Avian Influenza

Wild aquatic birds act as reservoirs for avian influenza viruses. Migratory movements facilitate long-distance

spread, posing risks to poultry industries and human health worldwide [16].

FUTURE PERSPECTIVES

Emerging technologies such as artificial intelligence, remote sensing, and environmental DNA analysis offer new opportunities for predicting and preventing zoonotic disease emergence. Strengthening global collaboration and investment in wildlife health research will be critical for long-term disease control.

CONCLUSION

Emerging infectious diseases in wildlife are driven by complex interactions among ecological, environmental, and socio-economic factors. Effective prevention and control require integrated strategies that combine ecological understanding, surveillance, conservation, and policy interventions. A strong One Health approach is essential to mitigate zoonotic risks while preserving biodiversity.

REFERENCES

1. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. *Nature*. 2008;451(7181):990–3.
2. Woolhouse ME, Gowtage-Sequeria S. Host range and emerging and re-emerging pathogens. *Emerg Infect Dis*. 2005;11(12):1842–7.
3. Luis AD, Hayman DTS, O’Shea TJ, Cryan PM, Gilbert AT, Pulliam JRC, et al. A comparison of bats and rodents as reservoirs of zoonotic viruses. *Proc Biol Sci*. 2013;280(1756):20122753.
4. Webster RG, Govorkova EA. Continuing challenges in influenza. *Ann N Y Acad Sci*. 2014;1323:115–39.
5. Hayman DTS. Bat tolerance to viral infections. *Nat Microbiol*. 2016;1:16050.
6. Patz JA, Daszak P, Tabor GM, Aguirre AA, Pearl M, Epstein J, et al. Unhealthy landscapes and disease emergence. *Environ Health Perspect*. 2004;112(10):1092–8.
7. Altizer S, Ostfeld RS, Johnson PTJ, Kutz S, Harvell CD. Climate change and infectious diseases. *Science*. 2013;341(6145):514–9.
8. Karesh WB, Cook RA, Bennett EL, Newcomb J. Wildlife trade and global disease emergence. *Emerg Infect Dis*. 2005;11(7):1000–2.
9. Ostfeld RS, Keesing F. Biodiversity and disease risk. *Conserv Biol*. 2000;14(3):722–8.
10. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic spillover. *Nat Rev Microbiol*. 2017;15(8):502–10.
11. Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife. *Science*. 2000;287(5452):443–9.
12. Atlas RM. One Health: its origins and future. *Curr Top Microbiol Immunol*. 2013;366:1–13.
13. Rupprecht CE, Hanlon CA, Hemachudha T. Rabies re-examined. *Lancet Infect Dis*. 2002;2(6):327–43.

14. Morse SS, Mazet JAK, Woolhouse M, Parrish CR, Carroll D, Karesh WB, et al. Prediction and prevention of pandemic zoonoses. *Lancet*. 2012;380(9857):1956–65.
15. Leroy EM, Epelboin A, Mondonge V, Pourrut X, Gonzalez JP, Muyembe-Tamfum JJ, et al. Human Ebola outbreak from fruit bats. *Vector Borne Zoonotic Dis*. 2009;9(6):723–8.
16. Olsen B, Munster VJ, Wallensten A, Waldenström J, Osterhaus ADME, Fouchier RAM. Global patterns of influenza A in wild birds. *Science*. 2006;312(5772):384–8.