



International Journal of Health Care and Biological Sciences

Review Article

Open Access

The osteoporosis treatment by gene therapy: a current approach and challenges

Riyakumari P.D.¹, Shreya M. S.², Trivedi Harsh^{3*}¹M.Pharm, Pharmaceutics, SDP College Gujarat 394110, India²B.Pharm Sigma Pharmacy College Bakrol, Vadodara Gujarat³B.Sc. (Biotechnology) Ganpat University Mehsana, India

Article History

Received on: 11-10-2021

Revised On : 25-10-2021

Accepted on : 03-12-2021

Keywords: Osteoporosis, Osteoporosis Management, Gene therapy.

DOI:

<https://doi.org/10.46795/ijhcbcs.v2i4.254>


Abstract

Osteoporosis is a worldwide disease characterized by reduction of bone mass and alteration of bone architecture resulting in increased bone fragility and increased fracture risk. Although it is seen in all age groups, gender, and races, it is more common in Caucasians (white race), older people, and women. With an aging population and longer life span, osteoporosis is increasingly becoming a global epidemic. Currently, it has been estimated that more than 200 million people are suffering from osteoporosis. Moreover, osteoporosis results in a decreased quality of life, increased disability-adjusted life span, and big financial burden to health insurance systems of countries that are responsible for the care of such patients. Therefore, increasing awareness in medical field, which, in turn, facilitates increase awareness of the normal populace, will be effective in preventing this epidemic.

This article is licensed under a Creative Commons Attribution-Non Commercial 4.0 International License. Copyright © 2021 Author(s) retain the copyright of this article.



*Corresponding Author

Trivedi Harsh

Email: sutharshreya343@gmail.com

Hosted by

saap.org.in

above 50 years age and 1 in 5 men above 50 years age are affected by osteoporotic fracture [3-5]. This shows that osteoporosis is a global healthcare burden.

Pathophysiology of The Disease

In osteoporosis, the bone mineral density (BMD) gets reduced, deterioration of bone microarchitecture takes place, and the amount of various transcription factors, growth factors and cytokines etc. in bone are altered. Imbalance between bone resorption and bone formation is the underlying mechanism in all cases of osteoporosis. In normal bone, matrix remodeling of bone is constant. Bone is resorbed by osteoclast cells, after which new bone is deposited by osteoblast cells. The three main mechanisms by which osteoporosis develop are (a) an inadequate peak bone mass (the skeleton develops insufficient mass and strength during growth), (b)

Introduction

Osteoporosis is a disease of bones that leads to an increased risk of fracture. It has been denoted a silent disease due to its character of occurring without symptomatic changes in the body. Worldwide estimates show that osteoporosis accounts for over 8.9 million fractures annually which turns out to be an osteoporosis related fracture every 3 seconds [1]. Osteoporosis affects 200 million women worldwide with approximately one-tenth of women aged 60 affected by osteoporosis [2]. It affects the aged people making them bedridden affecting their quality of life. Worldwide, 1 in 3 women

excessive bone resorption, and (c) inadequate formation of new bone during remodeling.

These occurs due to various hormonal level defects that lead to cascades of processes which cause increased bone resorption by osteoclasts and/or decreased bone generation by osteoblasts. Lack of estrogen (e.g. as a result of menopause) increases bone resorption, as well as decreasing the deposition of new bone that normally takes place in weight-bearing bones. In addition to estrogen, calcium metabolism plays a significant role in bone turnover, and deficiency of calcium and vitamin D leads to impaired bone deposition; in addition, the parathyroid glands react to low calcium levels by secreting parathyroid hormone (parathormone, PTH), which increases bone resorption to ensure sufficient calcium in the blood.

Main hormones that regulate bone metabolism are as follows: Decrease bone resorption: Calcitonin, estrogen, Increase bone resorption: parathormone (PTH), glucocorticoids, thyroid hormones, high dose vit.D, Increase bone formation: Growth hormone, vit. D metabolites, androgens, insulin, low dose PTH, Decrease bone formation: Glucocorticoids, Various growth factors, cytokines and transcription factors are involved in the pathogenesis of osteoporosis. These include RANK (receptor activator of nuclear factor $\kappa\beta$), RunX2 (Runt related factor X2), VEGF (vascular endothelial growth factor) [6], TNF (tumor necrosis factor), TGF (transforming growth factor), BMPs (bone morphogenetic proteins), OPG (osteoprotegerin), OTX (osterix). Many of these factors have been studied for their potential use in osteoporosis.

Osteoporosis Treatment and Its Management

Osteoporosis risk can be reduced with lifestyle changes and medication; in people with osteoporosis, treatment may involve both. Lifestyle change includes diet and exercise, and preventing falls. Medication includes supplemental calcium, vitamin D, calcitonin, bisphosphonates (zaledronic acid, ibandronate, etc), bone morphogenetic proteins (BMP-2 and -7) and several others [7]. Most of the therapies are long term therapies and require closer monitoring to avoid any adverse effects. Some of the demerits of the current therapeutics of osteoporosis are described in the Figure 1.1 below. Effectiveness of oral calcium and vitamin D supplementation has been evaluated extensively. The analyses show that calcium supplementation alone and vitamin D supplementation alone are not effective in

preventing fractures in osteoporotic patients as the combination thereof [8, 9]. Effect of intravenous calcium infusion has also been evaluated in osteoporotic women for treating osteoporosis, but it was found to be ineffective in altering bone calcium turnover in osteoporotic women. Loss of total body calcium was similar to that in untreated subjects with osteoporosis [10, 11].

Glucocorticoid-induced osteoporosis and osteoporosis related to aging are mainly outcome of reduced bone formation due to reduced number of osteoblasts. Moreover, calcium and vitamin D combination therapy has been found to be non-effective in preventing fractures in elderly (age >70 years) [12, 13]. An ideal way to prevent bone loss in such cases would be not only to reduce bone resorption, but also to promote bone formation. There is therefore an important need to develop therapeutic strategies capable of promoting bone formation in osteoporotic subjects.




Figure 1.1 Current treatment options of osteoporosis and their drawbacks

Gene Therapy as an Approach For Osteoporosis – Current Perspective

Gene delivery has been showing promising results for treatment of various diseases [14], [15], [16]. In the past decade various gene delivery approaches have been studied for the treatment of osteoporosis. Such gene delivery approaches particularly act either by inducing or one or other growth factors, cytokines, transcription factors, other mediators or their receptors that are implicated in osteoporosis. Advancements made in the treatment of osteoporosis with gene delivery are described below with brief review of various gene

delivery systems evaluated for osteoporosis treatment in animals (Figure 1.1).

Various cytokines, particularly interleukin-1 (IL-1) and tumor necrosis factor (TNF), have been strongly implicated in postmenopausal osteoporosis occurring due to estrogen deficiency. Both of these cytokines are powerful inducers of bone resorption. From this information, it follows that inhibiting the biological activities of IL-1 and TNF should reduce bone loss under conditions of estrogen deficiency [17], [18]. Genes encoding for IL-1 receptor antagonist (IL-1Ra) or soluble form of TNF receptors would ameliorate the osteoporotic bone loss by inhibiting osteoclastic activity [19, 20].

Intravenous delivery of human osteoprotegerin (hOPG) gene using viral vectors results in systemic circulation of the OPG which in turn inhibits osteoclastic activity. The mechanism involves the binding of OPG to RANKL (receptor activator of nuclear factor κ B ligand) which prevents the binding of latter to RANK. This in turn suppresses its ability to increase bone resorption by osteoclasts [21, 22]. LIM mineralization protein (LMP) which induces the bone mineralization and expression of various osteogenic genes, BMP-2, RunX2 (Runt related transcription factor X2), OSX (Osterix) etc., and thereby promotes the osteoblast differentiation. One study has also shown that it induces bone formation more efficiently than even BMP-2 [23].

Among all gene delivery approaches, delivery of genes of bone-morphogenetic proteins has been most extensively evaluated (Figure 1.1). Bone morphogenetic factors (BMPs), mainly BMP-2, BMP-4, BMP-6, BMP-7 and BMP-9, are other osteogenic proteins that have been studied for bone regeneration in fractured bone healing, osteoporosis and osteopenia [24]. Recombinant human bone morphogenetic protein-2 and -7 have been recently granted United States Food and Drug Administration approval for select clinical applications in bone repair (20, 24). These BMPs act primarily as differentiation factors, turning responsive mesenchymal cells into cartilage- and bone-forming cells [25]. While significant progress has been made in the delivery of recombinant osteogenic proteins to promote bone healing, the short half-life and instability of the protein requires the delivery of milligram quantities of factor or multiple dosages (20). So, delivery of genes encoding for various BMPs have been investigated in various studies (Figure 1.1). Various transcription factors and growth factors such as VEGF [26], RunX2, TGF etc. have also been found to enhance the effects of various BMPs. Among

various BMPs, BMP-9 has been shown to provide most robust and effective osteogenic activity in animal studies (Figure 1.1).

Conclusion

Osteoporosis is a skeletal disorder characterized by compromised bone strength leading to an increased risk of fracture. Osteoporosis is a common and silent disease until it is complicated by fractures that become common. It was estimated that 50% women and 20% of men over the age of 50 years will have an osteoporosis-related fracture in their remaining life. These fractures are responsible for lasting disability, impaired quality of life, and increased mortality, with enormous medical and heavy personnel burden on both the patient's and nation's economy. Therefore, the prevention, detection, and treatment of osteoporosis should be a mandate of primary healthcare providers.

Conflict of Interest

We declare that we have no conflict of interest.

References

1. Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*. 2006;17(12):1726-33.
2. WHO Technical report, WHO Press, Switzerland, 2007.
3. Melton LJ, 3rd, Atkinson EJ, O'Connor MK, O'Fallon WM, Riggs BL. Bone density and fracture risk in men. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 1998;13(12):1915-23.
4. Melton LJ, 3rd, Chrischilles EA, Cooper C, Lane AW, Riggs BL. Perspective. How many women have osteoporosis? *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*. 1992;7(9):1005-10.
5. Kanis JA, Johnell O, Oden A, Sembo I, Redlund-Johnell I, Dawson A, et al. Long-term risk of osteoporotic fracture in Malmo. *Osteoporosis international : a journal*

- established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA. 2000;11(8):669-74.
6. Bhatt P, Fnu G, Bhatia D, Shahid A, Sutariya V. Nanodelivery of Resveratrol-Loaded PLGA Nanoparticles for Age-Related Macular Degeneration. *AAPS PharmSciTech*. 2020;21(8):291.
 7. Vhora I, Lalani R, Bhatt P, Patil S, Patel H, Patel V, et al. Colloidally Stable Small Unilamellar Stearyl Amine Lipoplexes for Effective BMP-9 Gene Delivery to Stem Cells for Osteogenic Differentiation. *AAPS PharmSciTech* 2018; 19:3550–60.
 8. Garcia Vadillo JA. [Are calcium and vitamin D supplements for everyone?: Pro]. *Reumatologia clinica*. 2011;7 Suppl 2:S34-9.
 9. Nowson CA. Prevention of fractures in older people with calcium and vitamin D. *Nutrients*. 2010;2(9):975-84.
 10. Dudl RJ, Ensinnck JW, Baylink D, Chesnut CH, 3rd, Sherrard D, Nelp WB, et al. Evaluation of intravenous calcium as therapy for osteoporosis. *The American journal of medicine*. 1973;55(5):631-7.
 11. Jensen H, Christiansen C, Munck O, Toft H. Treatment of osteoporosis with calcium infusions. An osteodensitometric study. *Scandinavian journal of clinical and laboratory investigation*. 1973;32(1):93-6.
 12. Grant AM, Avenell A, Campbell MK, McDonald AM, MacLennan GS, McPherson GC, et al. Oral vitamin D3 and calcium for secondary prevention of low-trauma fractures in elderly people (Randomised Evaluation of Calcium Or vitamin D, RECORD): a randomised placebo-controlled trial. *Lancet*. 2005;365(9471):1621-8.
 13. Porthouse J, Cockayne S, King C, Saxon L, Steele E, Aspray T, et al. Randomised controlled trial of calcium and supplementation with cholecalciferol (vitamin D3) for prevention of fractures in primary care. *Bmj*. 2005;330(7498):1003.
 14. Patil S, Bhatt P, Lalani R, Amrutiya J, Vhora I, Kolte A, et al. Low molecular weight chitosan-protamine conjugate for siRNA delivery with enhanced stability and transfection efficiency. *RSC Advances*. 2016;6(112):110951-63.
 15. Lalani R, Misra A, Amrutiya J, Patel H, Bhatt P, Patil SK. Approaches and Recent Trends in Gene Delivery for Treatment of Atherosclerosis. *Recent patents on drug delivery & formulation*. 2016;10(2):141-55.
 16. Patil S, Lalani R, Bhatt P, Vhora I, Patel V, Patel H, et al. Hydroxyethyl substituted linear polyethylenimine for safe and efficient delivery of siRNA therapeutics. *RSC Advances*. 2018;8(62):35461-73.
 17. Vhora I, Patil S, Bhatt P, Gandhi R, Baradia D, Misra A. Receptor-targeted drug delivery: current perspective and challenges. *Therapeutic delivery*. 2014;5(9):1007-24.
 18. Vhora I, Lalani R, Bhatt P, Patil S, Misra A. Lipid-nucleic acid nanoparticles of novel ionizable lipids for systemic BMP-9 gene delivery to bone-marrow mesenchymal stem cells for osteoinduction. *International Journal of Pharmaceutics*. 2019;563:324-36.
 19. Baltzer AW, Whalen JD, Wooley P, Latterman C, Truchan LM, Robbins PD, et al. Gene therapy for osteoporosis: evaluation in a murine ovariectomy model. *Gene therapy*. 2001;8(23):1770-6.
 20. Kofron MD, Laurencin CT. Bone tissue engineering by gene delivery. *Advanced drug delivery reviews*. 2006;58(4):555-76.
 21. Kostenuik PJ, Bolon B, Morony S, Daris M, Geng Z, Carter C, et al. Gene therapy with human recombinant osteoprotegerin reverses established osteopenia in ovariectomized mice. *Bone*. 2004;34(4):656-64.
 22. Ulrich-Vinther M, Carmody EE, Goater JJ, K Sb, O'Keefe RJ, Schwarz EM. Recombinant adeno-associated virus-mediated osteoprotegerin gene therapy inhibits wear debris-induced osteolysis. *The Journal of bone and joint surgery American volume*. 2002;84-A(8):1405-12.
 23. Pola E, Gao W, Zhou Y, Pola R, Lattanzi W, Sfeir C, et al. Efficient bone formation by gene transfer of human LIM mineralization protein-3. *Gene therapy*. 2004;11(8):683-93.
 24. Gautschi OP, Frey SP, Zellweger R. Bone morphogenetic proteins in clinical applications. *ANZ journal of surgery*. 2007;77(8):626-31.

25. Ebara S, Nakayama K. Mechanism for the action of bone morphogenetic proteins and regulation of their activity. *Spine*. 2002;27(16 Suppl 1):S10-5.
26. Bhatt P, Narvekar P, Lalani R, Chougule MB, Pathak Y, Sutariya V. An in vitro Assessment of Thermo-Reversible Gel Formulation Containing Sunitinib Nanoparticles for Neovascular Age-Related Macular Degeneration. *AAPS PharmSciTech*. 2019;20(7):281.