



# International Journal of Health Care and Biological Sciences

Review Article

Open Access

Measures of disease frequency and prevalence of drug therapy related problems on basis of prospective observational study in rural hospitalization patients in Nellore

A. Sushma<sup>1</sup>, Ch. Venkata Suseela<sup>1</sup>, B. Hepsi<sup>1</sup>, P. Mujib Ahmed<sup>1</sup>, S. Venkata Reddy<sup>1</sup>, C.Kiran Kumar<sup>2</sup>, P. Venkatesh<sup>3</sup>

<sup>1</sup> B.Pharmacy final year student, Jagan's Institute of Pharmaceutical Sciences, Nellore

<sup>2</sup> Associate Professor, Dept.of Pharmacy practice, Jagan's Institute of Pharmaceutical Science, Nellore

<sup>3</sup> Principal, Jagan's Institute of Pharmaceutical Science, Nellore

## Article History

Received on: 28-04-2021

Revised on: 10-05-2021

Accepted on: 25-06-2022

**Keywords:** Drug-related problems, Phenytoin, Nifedipine.

DOI:

<https://doi.org/10.46795/ijhcbcs.v3i2.330>



## Abstract

**Aim and objectives:** The study is aimed to assess the frequency and prevalence of drug therapy-related problems in patients after hospitalization in Jayabharath Hospital, Nellore. **Specific objectives:** To estimate the opportunities for pharmacists to provide pharmaceutical care. A probabilistic convenient sampling was done to estimate the frequency and prevalence of drug therapy-related problems, patients who are aged above 18 years of either sex were included. Pregnant patients, and those who are having an unwillingness to participate in study were excluded. **Methodology:** This cross-sectional study was conducted between March 2022 to June 2022 on patients who were discharged home from Jayabharath Hospital, Nellore. Most of our study population were within the age group of 30 to 45 years (33.33 %) followed by 60 to 75 years (25.00 %). among the present study population most of them were married (83.3 %) and illiterates (62.5 %). **Results and Discussion:** Nearly 50 drugs are responsible for drug-related problems in the present study among those Phenytoin was found to be the leading drug in relation to the development of drug therapy-related problems, Whereas Nifedipine was found to be least problem-producing the drug in the present study population. **Conclusion:** Overall we found various kinds of drug therapy-related problems, which implies there is a need for continuity of care even after discharge from the hospital.

This article is licensed under a Creative Commons Attribution-Non-commercial 4.0 International License. Copyright © 2022 Author(s) retain the copyright of this article.



## \*Corresponding Author

C. Kiran Kumar

Hosted by

saap.org.in

## Introduction

Pharmaceutical care

Pharmaceutical care is the responsible provision of drug therapy for the purpose of achieving definite outcomes that improve a patient's quality of life. It involves the process through which a pharmacist cooperates with a patient and other professionals in designing, implementing, and monitoring a therapeutic plan that

will produce specific therapeutic outcomes for the patient. This in turn involves three major functions:

- Identifying potential and actual drug-related problems;
- Resolving actual drug-related problems; and
- Preventing drug-related problems.

Pharmaceutical care is a necessary element of health care and should be integrated with other elements. Pharmaceutical care is, however, provided for the direct benefit of the patient, and the pharmacist is responsible directly to the patient for the quality of that care. The fundamental relationship in pharmaceutical care is a mutually beneficial exchange in which the patient grants authority to the provider, and the provider gives competence and commitment (accept responsibility) to the patient. The fundamental goals, processes, and relationships of pharmaceutical care exist regardless of the practice setting [1].

Principles of Practice for Pharmaceutical Care: - A professional relationship: Interaction between the pharmacist and the patient must occur to assure that a relationship based upon caring, trust, open communication, cooperation, and mutual decision making is established and maintained. In this relationship, the pharmacist holds the patient's welfare paramount, maintains an appropriate attitude of caring for the patient's welfare, and uses all his/her professional knowledge and skills on the patient's behalf [2].

Maintenance and documentation of patient information: Pharmacists must collect and/or generate subjective and objective information regarding the patient's general health and activity status, past medical history, medication history, social history, diet and exercise history, history of present illness, and economic situation. Sources of information may include, but are not limited to, the patient, medical charts and reports, direct patient interview by pharmacist, obtaining information from the patient's family or caregiver, insurer, and other healthcare providers including physicians, nurses, mid-level practitioners and other pharmacists [2].

Evaluation of patient medical information and development of drug therapy plan:

Based upon a thorough understanding of the patient and his/her condition or disease and its treatment, the pharmacist must, with the patient and with the patient's other healthcare providers as necessary, develop an outcomes-oriented drug therapy plan. The plan may have various components which address each of the

patient's diseases or conditions. In designing the plan, the pharmacist must carefully consider the psychosocial aspects of the disease as well as the potential relationship between the cost and/or complexity of therapy and patient adherence. As one of the patient's advocates, the pharmacist assures the coordination of drug therapy with the patient's other healthcare providers and the patient [2].

#### Study Procedure

- A cross-sectional observational studies was conducted on March 2022 and June 2022 in Jayabharath Hospital, Nellore, India. This study was approved by Institution Review Boards of Jayabharat Hospital, Nellore, India. All the subjects data obtained from medical record department of Jayabharath Hospital, Nellore.
- Subjects without pregnancy ,on any gender who are suffering from adverse reactions.who are under use of medication ,drug duplication ,drug –drug interactions, therapy failure ,practical problems of administrating dose ,missing doses were included in the study. Totally 100 patients with above reasons were included in the study remaining are excluded.

#### Statistical Analysis

- Beer's criteria was used to estimate the priority and severity of drug related problems and to quantify the drug therapy related problems. Drug therapy related problems are estimated by using medication appropriate index.
- Descriptive statistics were used to analyze the data.
- MAI uses 10 criteria for each medication a patient is taking, which assess medication indication, effectiveness, dosage, directions, drug-drug interactions, drug-disease interactions, expense, practicality, duplication, and treatment duration.

#### Results

The demographic details are presented in Table 01. It shows that most of the people in the present study were within the age group of 30 to 45years (33.33 %) followed by 60 to 75years (25.00%). Among the present study population most of them were married (86.11%), salaried (33.33%), illiterates (97.64 %), less than medications taken (58.33%) and hospitalized in past

year patients are (81.94%) The present study comprises of more than 40 % of patients with more than or equal to five medications and also around 80% of patients are having a history of previous hospitalization.

**Table 01: Demographic characteristics of the study subjects**

S. No	Characteristics details	Number of subjects N=72 (%)
1	<b>Age:</b>	
	18 – 30 years	10(13.88)
	30 – 45 years	30(33.33)
	45 – 60 years	14(19.44)
2	60 – 75 years	18(25.00).
	<b>Marital status:</b>	
	Married	62(86.11)
	Unmarried	10(13.88)
3	Divorced	00(00.00)
	Widowed	00(00.00)
	<b>Occupation:</b>	
	Business	18(25.00)
	Farmer	16(19.44)
4	salaried	24(33.33)
	Daily wager	12(13.88)
	Student	2(2.16)
	<b>Educational status:</b>	
	Illiterate	68(97.64)
5	Primary school	0
	High school	1(1.38)
	Inter	3(4.16)
	Degree	0
5	No. of medications:	
	Less than five	42(58.33)
	More than five	30(41.66)

Various kinds of drug therapy related problems are described in Table II and their Frequency and prevalence were detailed in Table 03. We found drug – drug interactions are the most frequent kind of drug related problems, which included only one interaction (37.5%) and more than one interaction per patient (62.5%). Additive toxicity was found to be second highest (25 %) drug related problems in the present study subjects, and we do not found any kind of drug related problems related to Duration of therapy.

**Table 02: Classification of drug therapy related problems**

Types of drug-related problems
<ol style="list-style-type: none"> <li>1. Uncertainty and lack of knowledge.</li> <li>2. Under use of medication.</li> <li>3. Over use of medication.</li> <li>4. Other dosage problems.</li> <li>5. Drug duplication.</li> <li>6. Drug - Drug interactions.</li> <li>7. Therapy failure.</li> <li>8. Side effect.</li> <li>9. Difficulty in swallowing / difficulty in dosage.</li> <li>10. Practical problems of administering dose.</li> <li>11. Language difficulty / understanding disability.</li> <li>12. Prescribing error such as incorrect or omitted data on prescribing drug.</li> </ol>

**Table 03: Frequency and prevalence of drug related problems:**

S.No	Drug-related problem category	No. of Patients withat least one DRP (%)	No. of Patients with more than one DRP (%)
1	Therapeutic Duplication	8(11.11)	4(5.55)
2	Dosage form problems	2(2.77)	1(1.38)
3	Untreated indication	6(8.33)	2(2.77)
4	Duration of therapy	0(0.00)	0(0.00)
5	Drug - Drug interactions	27(37.50)	45(62.50)
6	Additive toxicity	12(16.66)	18(25.00)
7	Non adherence	2(2.77)	1(1.38)
8	Treatment of noindication	1(1.38)	10(13.99)
9	Sub therapeutic dose	0(0.00)	1(1.38)
10	Improper drug selection	8(11.11)	3(4.16)

According to Beer's criteria Drug therapy problems were specified in Table IV. Which explains about the outcomes of the problem whether it is clinically significant or not and it also states about their priority and severity.

Table IV explains Drug – Drug interactions (140.44) and adverse drugs reactions (122.33) are the most commonly found Drug related problems which is clinically significant and even they considered as prime criteria to resolve the issue according to Beer's criteria. Social history, recreational drug use and adverse drug reactions makes the drug therapy related problems to be more severe.

**Table 05: Drug therapy problems according to Beers criteria**

S.No	Type of drug therapy problems	Outcome of problem		Priority of problem			Severity of problem		
		Clinical	Non-clinical	Primary	Secondary	Tertiary	Minor	Moderate	Major
1	Untreated indication	8(11.33)	2(2.77)	2(2.77)	3(4.16)	6(8.33)	6(8.33)	3(4.16)	2(2.77)
2	Improper drug selection	10(13.99)	4(5.55)	4(5.55)	2(2.77)	8(11.33)	8(11.33)	2(2.77)	4(5.55)
3	Sub therapeutic dose	1(1.38)	0	0	1(1.38)	0	0	1(1.38)	0
4	Over dose	1(1.38)	0	0	0	1(1.38)	1(1.38)	0	0
5	Adverse drug reaction	86(119.55)	2(2.77)	21(29.16)	67(93.05)	1(1.38)	1(1.38)	67(93.05)	21(29.16)
6	Drug drug interaction	141(139.5)	1(1.38)	42(58.33)	81(112.5)	18(25.0)	18(25.0)	81(112.5)	42(58.33)
7	Non adherence	1(1.38)	3(4.16)	0	2(2.77)	2(2.77)	2(2.77)	2(2.77)	0
8	No indication	12(16.66)	2(2.77)	3(4.16)	4(5.55)	5(6.94)	5(6.94)	4(5.55)	3(4.16)
9	Social, recreational drug use	30(41.55)	0	27(37.5)	3(4.16)	0	0	3(4.16)	27(37.5)
10	Patient knowledge on drug therapy	0	12(16.66)	0	0	12(16.66)	12(16.66)	0	0

Table 05. Shows various drugs responsible for the drug related problems in the present study population. Nearly 50 drugs are responsible for drug related problems in the present study among those Phenytoin was found to be leading drug in relation to development of drug therapy related problems, Whereas Nifedipine was found to be least problem producing drug in present study population.

**Table V. Drugs implicated for problems**

Drugs	Number = (%)
Amoxicillin	10(22.72)
Amikacin	5(11.36)
Ceftriaxone	7(15.90)
Doxycycline	4(9.09)
Erythromycin	3(6.81)
Vancomycin	2(4.54)
Azithromycin	8(18.18)
Ciprofloxacin	4(9.09)
Furosemide	7(15.90)

Spiranolactone	3(6.81)
Digoxin	2(4.54)
Atorvastatin	5(11.36)
Amlodipine	6(13.63)
Simvastatin	2(4.54)
Nifedipine	1(2.27)
Lisinopril	3(6.81)
Ramipril	9(20.54)
Propranolol	8(18.18)
Atenolol	6(13.63)
Rifampicin	12(27.27)
Isoniazid	11(25.0)
Ranitidine	16(36.36)
Phenytoin	21(47.72)
Carbamazepine	12(27.27)
Valproic acid	7(15.90)
Diazepam	18(40.90)
Fluoxetine	12(27.27)
Clomipramine	8(18.18)
Trazadone	6(13.63)
Amitryptalin	2(4.54)
Tramadol	5(11.36)
Salbutamol	6(13.63)
Terbutaline	4(9.09)
Prednisolone	9(20.54)
Aspirin	8(18.18)
Celecoxib	6(13.63)
Metformin	9(20.54)
Glimepramide	12(27.27)
Omeprazole	8(18.18)
Paracetamol	16(36.36)
Ibuprofen	6(13.63)
Diclofenac	5(11.63)
Cetirizine	6(13.63)

## Discussion

The present study shows that DRPs among patients discharged from the hospital 93.7% of patients in this study reported DRPs. Although most DRPs do not seem to have direct clinical complications, they could lead to patient education nonadherence which is another drug therapy related problem which affects the patient psychology to avoid the medications in near future.<sup>45</sup> The present study results shows that there is a huge number of drug related problems which implies that there is a need of care by health care professionals even after hospitalization, preferably by pharmacist to prevent or resolve those drug related problems, this

confirms discharged patients are an important target group for pharmacy interventions.

In a study the medication review performed by the pharmacists resulted in 476 identified DRPs of which the most common was adverse drug reaction (119.8%), commonly caused by a dose that was too high and not adjusted to the patient's physical status and elimination capacity. The second most common DRP was need for additional drug therapy and unnecessary drug therapy (86%).<sup>44</sup> But in our study we observed Drug - Drug interactions and adverse drug reactions as most common drug therapy problems (219.15).

Several authors have reported that a follow-up home visit by a pharmacist had no benefit for patients who had been discharged from hospital. It is difficult to reconcile those data with data illustrating a benefit of medication related support during the discharge period, but those trials were less focused on identifying and resolving drug-related problems or that the support was not received early enough after the time of discharge.

However, these contradictory data represent an opportunity for further investigation into what sorts of best support for patients with their medication therapy can be provided by pharmacist after discharge from acute care.

Varying distributions of DRPs in relation to different types of drugs have been reported in previous studies. But in the present study gastrointestinal drugs and central nervous system drugs were highly involved. Our finding shows that the number of new drugs, continued drugs and changes in dosage at discharge are independently associated with the occurrence of DRPs which is similar to the results of other hospital based studies.

In a previous study, 12% of all visits to the emergency department were due to adverse drug events; 68% of these adverse events were considered preventable in both of the study. A consultation with a pharmacist after discharge from hospital presents an opportunity for these errors to be identified and recommendations to be made. It also provides an opportunity to improve patients' knowledge of their medications.

## Limitations

The present study comprises of less number of study sample and more over it is a preliminary study to evaluate drug related problems (DRP) to interventions were not provided, so further the present study can be continued after the implementation of pharmacist intervention to evaluate pharmaceutical care.

## Conclusion

Overall we found various kinds of drug therapy related problems, which implies there is a need of continuity of care even after discharge from hospital. Preferably pharmacist's intervention will be useful to reduce these problems, which clearly shows a greater opportunity for pharmacist to provide pharmaceutical care to the patients those who are discharged from the hospital.

## Funding

No Funding

## Conflict of Interest

The authors have declared no conflict of interest.

## Author Contribution

All authors contributed equally.

## References

1. Helper DD, Strand LM, et al., Opportunities and Responsibilities in Pharmaceutical Care. *American journal of hospital pharmacy*. 1989; 47(3), 7-15.
2. American public health association Pharmaceutical Care Guidelines Advisory Committee, [www.marketresearch.com/product/display.asp?productid=1327146&](http://www.marketresearch.com/product/display.asp?productid=1327146&) APhA Board of Trustees, 1995.
3. The pharmaceutical market: Peru. Espicom Healthcare Intelligence. 2006. [www.marketresearch.com/product/display.asp?productid=1327146&](http://www.marketresearch.com/product/display.asp?productid=1327146&) (accessed 2007).
4. FIP Statement of Policy on Collaborative Pharmacy Practice. Statements and guide lines of pharmaceutical care. 1995, 20-21.
5. Mikeal RL, Brown TR, Lazarus HL, et al., Quality of Pharmaceutical care in hospitals. *American journal of hospital pharmacy*. 1975; 32(5): 567-574.
6. Brodie DC. Drug Use Control: keystone to pharmaceutical service. *Drug Intelligence and Clinical Pharmacy*. 1967; 1(1):63-65.
7. Bond CA, Raehl CL, Franke T. Interrelationships among mortality rates, drug costs, total costs of care and length of stay in United States hospital, summary and recommendations for clinical pharmacy services and staffing. *Pharmacotherapy* 2001; 21(4):129-141.
8. Parthasarathi G, Ramesh M, Hansen KN, Nagavi BG. Clinical pharmacy in a South Indian teaching hospital. *The annals of Pharmacotherapy* 2002; 36(5): 927-932.
9. Geer MI, Mir JI, Koul PA. Inauguration of Pharmaceutical Care Clinic for patient safety and therapeutic efficacy for chronic diseases management, drug information and pharmacoconomics. *International journal of community pharmacy* 2009; 2(3): 1-4.
10. Kannan G, Janardhan V, Rani V, et al., Pharmaceutical care in the general medicine ward of a tertiary care hospital in South India. *Journal of Pharmacy Res* 2011; 4(5): 1467- 1469.
11. PCNE. The PCNE Classification V 5.01 2006 29 May 2006 [cited 20 Sep 2009]. Available from: <http://www.pcne.org/dokumenter/DRP/PCNE%20classification%20V5.01.pdf>.
12. AbuRuz SM, Bulatova NR, Yousef AM. Validation of a comprehensive classification tool for treatment-related problems. *Pharma world science journal*. 2006; 28(4):222-232.
13. Westerlund LT, Almarsdottir AB, Melander A. Drug-related problems and pharmacy interventions in community practice. *International journal of pharmacy*. 1999; 7(11): 40- 50.
14. Rawlins MD, Thompson JW. Mechanisms of adverse drug reactions In: Davies DM, editor. *Textbook of adverse drug reactions*. Oxford University Press, 1991: 18-45.
15. Inman WHW, editor. *Monitoring for drug safety*. 2nd ed. Lancaster: MTP Press, 1986. [www.who.com](http://www.who.com) [www.who-umc.org/graphics/25257.pdf](http://www.who-umc.org/graphics/25257.pdf)
16. Royer RJ. Mechanism of action of adverse drug reactions: an overview *Pharma epidemiology Drug Saftey* 1997; 3: 43-S50.